

Tyrosinase Inhibitors Isolated from the Edible Brown Alga *Ecklonia stolonifera*

Hye Sook Kang, Hyung Rak Kim, Dae Seok Byun, Byeng Wha Son¹, Taek Jeong Nam, and Jae Sue Choi
Faculty of Food Science and Biotechnology and ¹Department of Chemistry, Pukyong National University, Busan
608-737, Korea

(Received September 10, 2004)

Extracts from seventeen seaweeds were determined for tyrosinase inhibitory activity using mushroom tyrosinase with L-tyrosine as a substrate. Only one of them, *Ecklonia stolonifera* OKAMURA (Laminariaceae) belonging to brown algae, showed high tyrosinase inhibitory activity. Bioassay-guided fractionation of the active ethyl acetate (EtOAc) soluble fraction from the methanolic extract of *E. stolonifera*, led us to the isolation of phloroglucinol derivatives [phloroglucinol (1), eckstolonol (2), eckol (3), phlorofucofuroeckol A (4), and dieckol (5)]. Compounds 1-5 were found to inhibit the oxidation of L-tyrosine catalyzed by mushroom tyrosinase with IC_{50} values of 92.8, 126, 33.2, 177, and 2.16 $\mu\text{g/mL}$, respectively. It was compared with those of kojic acid and arbutin, well-known tyrosinase inhibitors, with IC_{50} values of 6.32 and 112 $\mu\text{g/mL}$, respectively. The inhibitory kinetics analyzed from Lineweaver-Burk plots, showed compounds 1 and 2 to be competitive inhibitors with K_i of 2.3×10^{-4} and 3.1×10^{-4} M, and compounds 3-5 to be noncompetitive inhibitors with K_i of 1.9×10^{-5} , 1.4×10^{-3} and 1.5×10^{-5} M, respectively. This work showed that phloroglucinol derivatives, natural compounds found in brown algae, could be involved in the control of pigmentation in plants and other organisms through inhibition of tyrosinase activity using L-tyrosine as a substrate.

Key words: *Ecklonia stolonifera*, Tyrosinase inhibitors, Phloroglucinol derivatives, Marine algae, Kinetics

INTRODUCTION

Tyrosinase (EC 1.14.18.1) is a multifunctional, copper-containing oxidase that catalyzes three distinct reactions of melanin synthesis, the hydroxylation of tyrosine to 3,4-dihydroxyphenylalanine (DOPA), the oxidation of DOPA to dopaquinone and the conversion of 5,6-dihydroxyindole to melanochrome (Mason, 1956; Wilcox *et al.*, 1985). Tyrosinase also known as polyphenol oxidase (PPO) is ubiquitously distributed from microorganisms to animals and plants (Mayer, 1987; Whitaker, 1995). The enzyme induces the browning of some fruits, vegetables, and crustaceans and causes a significant discount in their nutritional and market values. In addition, tyrosinase is a key enzyme in the insect molting process, so that its inhibitors might ultimately provide clues to control insect pests (Andersen,

1979). The enzymatic oxidation of L-tyrosine to melanin is of considerable importance because melanin has many functions, and alterations in melanin synthesis occur in many disease states. Recently, tyrosinase inhibitors have become increasingly important in cosmetic and medical products in relation to hyperpigmentation (Kim *et al.*, 2002; Pérez-Bernal *et al.*, 2000). Furthermore, it has been reported that tyrosinase could be central to dopamine neurotoxicity as well as contributing to the neurodegeneration associated with Parkinson's disease (Xu *et al.*, 1997).

Ecklonia stolonifera OKAMURA (Laminariaceae), a perennial brown alga growing in a water depth of 2-10 m, is distributed in Korea and Japan, and is commonly used as a foodstuff along with *Laminaria japonica* and *Undaria pinnatifida*. Phloroglucinol and its derivatives (Taniguchi *et al.*, 1991), and ecklonialactones (Kurata *et al.*, 1989; 1993) have been isolated from *E. stolonifera*. This alga was found to have antioxidant (Choi *et al.*, 1993; Lee *et al.*, 1996), antimutagenic activity (Lee *et al.*, 1996; 1998; Han *et al.*, 2000), and feeding-deterrent effect (Taniguchi *et al.*, 1991). However, the inhibitory effect on the tyrosinase

Correspondence to: Jae Sue Choi, Faculty of Food Science and Biotechnology, Pukyong National University, Busan 608-737, Korea
Tel: 82-51-620-6335, Fax: 82-51-620-6330
E-mail: choijs@pknu.ac.kr

activity was not investigated. We report the phloroglucinol derivatives with tyrosinase inhibitory activity from *E. stolonifera*, and the inhibitory kinetics of the components on this enzyme activity.

MATERIALS AND METHODS

Plant material

Leafy thalli of the *E. stolonifera* were collected at Gijang-gun in Busan, South Korea in February 2000 and authenticated by Prof. H. G. Kim of the Faculty of Marine Bioscience and Technology, Kangnung National University. A voucher specimen (no. 20000228) has been deposited in the author's laboratory (JS Choi).

Chemicals and reagents

Column chromatography was done with silica gel 60 (70~230 mesh, Merck, Germany), RP-18 Lichroprep (Merck, Germany), and Sephadex LH-20 (Sigma, St. Louis, MO). TLC was carried out on a precoated Merck Kieselgel 60 F₂₅₄ plate (0.25 mm), and 50% H₂SO₄ was used as spray reagent. L-Tyrosine was purchased from Janssen Chimica (Geel Belgium). Arbutin, kojic acid and mushroom tyrosinase (EC 1.14.18.1) were obtained from Sigma Chemical Co. K₂HPO₄ was obtained from Junsei Chemical Co. Ltd. (Tokyo, Japan), and KH₂PO₄ from Yakuri Pure Chemicals Co. Ltd. (Osaka, Japan). All the solvents for column chromatography were of reagent grade from commercial sources.

Isolation of active compounds

The lyophilized powder (3 kg) was refluxed with MeOH (3×9 L) for 3 h. The extract (700 g) was suspended in water and partitioned with *n*-hexane (27.9 g), CH₂Cl₂ (25.6 g), EtOAc (25.0 g), *n*-BuOH (99.6 g), in sequence. The EtOAc fraction (25.0 g) was applied to a silica gel (Merck, 70~230 mesh, 4×80 cm, 800 g) column. The column was eluted using mixtures of EtOAc/MeOH under stepwise gradient conditions (50:1~5:1) to yield the 10 subfractions (F1~F10), i.e., F1~F3; EtOAc/MeOH, 50:1 (5 L), F4~F6; EtOAc/MeOH, 10:1 (5 L), F7~F8; EtOAc/MeOH, 5:1 (5 L), and F9~F10; EtOAc/MeOH, 2:1 (2 L). The F1 (3.44 g, IC₅₀ = 20 µg/mL) was subjected to a silica gel column chromatography (70~230 mesh, 3×70 cm, 250 g, *n*-hexane/EtOAc, 1:1) to get the 11 subfractions (F1-1~F1-11). Compound 1 (98 mg) was obtained from the RP-18 column chromatography (20% MeOH~100% MeOH, gradient) of F1-4 (257 mg). Compounds 2 (60 mg) and 3 (135 mg) were obtained from the RP-18 column chromatography (20% MeOH~100% MeOH, gradient) of F1-5 (1.01 g). Compounds 4 (57 mg) and 5 (87 mg) in F1-6 (945 mg) were obtained from the RP-18 column chromatography using a 20% MeOH~100% MeOH gradient, then finally purified by

Sephadex LH-20 column chromatography with MeOH as a solvent.

The isolated compounds were identified as phloroglucinol (1), eckstolonol (2), eckol (3), phlorofucofuroeckol A (4), and dieckol (5) on the basis of chemical and physico-chemical evidences, and compared the spectral data with those in the literatures reported previously (Fukuyama *et al.*, 1985; 1989; 1990; Nakamura *et al.*, 1996; Kang *et al.*, 2003).

Enzyme assay

Tyrosinase activity using L-tyrosine as a substrate was spectrophotometrically determined by a method (No *et al.*, 1999) described previously with a little modification. Ten µL of each sample solution with different concentrations (1~500 µg/mL) and 20 µL of mushroom tyrosinase (1000 units/mL) in a 50 mM phosphate buffer (pH 6.5) were added to 170 µL of an assay mixture containing with the ratio 10:10:9 of 1 mM L-tyrosine solution, 50 mM potassium phosphate buffer (pH 6.5) and distilled water in a 96-well microplate. One unit (U) of enzyme activity was defined as the amount of enzyme increasing 0.001 absorbance at 280 nm per min, in 3 mL reaction mix containing L-tyrosine at pH 6.5 at 25°C. The samples dissolved in DMSO were diluted to 30 times with distilled water before experiment. After incubation of the reaction mixture at 25 °C for 30 min, the absorbance of the mixture was determined at 490 nm ($\epsilon = 3.3 \times 10^3 \text{ M}^{-1}\text{cm}^{-1}$) in a microplate reader (VERSA max, Molecular Device, CA). The extent of inhibition by addition of samples is expressed as a concentration necessary for 50% inhibition (IC₅₀). The percent inhibition of tyrosinase activity was calculated by the following equation: the % Inhibition = $[(1 - (A_a - A_b)/A_c) \times 100]$ where A_a is absorbance at 490 nm with test sample and enzyme, A_b is absorbance at 490 nm with test sample and without enzyme, and A_c is absorbance at 490 nm with enzyme and without test sample.

The preincubation mixture consisted of 60 µL of 50 mM phosphate buffer (pH 6.5), 50 µL of water, 10 µL of the samples solution, and 20 µL of the aqueous solution of the mushroom tyrosinase (1000 U/mL). The mixture was preincubated at 25°C for 5 min. Then, 60 µL of 1 mM L-tyrosine was added, and the reaction was monitored at 490 nm after 2 min incubation.

Kinetic analysis

The reaction mixture consisted of four different concentrations of L-tyrosine (0.5 to 2 mM) as a substrate and mushroom tyrosinase in 50 mM potassium phosphate buffer. Each sample of several concentrations was added to the reaction mixture, respectively. Michaelis constant (K_m) and maximal velocity (V_{max}) of the tyrosinase were determined by Lineweaver-Burk plots. The velocity equation

for the competitive inhibition in reciprocal form is: $1/V = K_m / V_{max} (1 + [I] / K_i) 1/[S] + 1/V_{max}$. Inhibition constants (K_i) of the competitive inhibitors were calculated by the following equation: $Km_{app} = K_m [1 + ([I] / K_i)]$ where Km_{app} is the apparent K_m in the presence of any inhibitor concentration. The reciprocal equation for the noncompetitive inhibition is: $1/V = K_m / V_{max} (1 + [I] / K_i) 1/[S] + 1/V_{max} (1 + [I] / K_i)$. K_i of the noncompetitive inhibitors were calculated by the following equation: $1/V_{max_{app}} = (1 + [I] / K_i) / V_{max}$ where $V_{max_{app}}$ is the apparent V_{max} in the presence of any inhibitor concentration.

RESULTS AND DISCUSSION

To discover the sources with tyrosinase inhibitory activity in marine algae, 17 seaweed extracts *i.e.* 2 of Chlorophyta, 5 of Phaeophyta, and 10 of Rhodophyta, were screened for their inhibitory effect on mushroom tyrosinase activity using L-tyrosine as a substrate (Table I). Only one of them, *Ecklonia stolonifera* OKAMURA (Laminariaceae), which belongs to brown algae, showed an inhibitory activity with an IC_{50} value of 345 $\mu\text{g/mL}$. The others showed an inhibitory activity of below fifty percent at the highest concentration of 500 $\mu\text{g/mL}$.

In our ongoing study to identify the active principles, we also evaluated the solvent soluble fractions including *n*-hexane, CH_2Cl_2 , EtOAc, *n*-BuOH, as well as the H_2O layer derived from *E. stolonifera*. As summarized in Table

Table I. Inhibitory activities of seaweeds on mushroom tyrosinase

Seaweeds		IC_{50}^a ($\mu\text{g/mL}$)
Chlorophyta	<i>Enteromorpha crinita</i>	>500
	<i>Ulva pertusa</i>	>500
Phaeophyta	<i>Ecklonia stolonifera</i>	345
	<i>Hizikia fusiformis</i>	>500
	<i>Sargassum ringgoldianum</i>	>500
	<i>Sargassum miyabei</i>	>500
	<i>Sargassum thunbergii</i>	>500
Rhodophyta	<i>Acrosorium flabellatum</i>	>500
	<i>Carpopeltis comea</i>	>500
	<i>Chondrus crispus</i>	>500
	<i>Gelidium amansii</i>	>500
	<i>Gigatina tenella</i>	>500
	<i>Gracilaria verrucosa</i>	>500
	<i>Grateloupia filicina</i>	>500
	<i>Pachymeniopsis elliptica</i>	>500
<i>Rhodymenia intricate</i>	>500	
<i>Symphocladia latiuscula</i>	>500	
Control	Kojic acid	4.57

^aInhibitory activity was expressed as the mean of 50% inhibitory concentrations of triplicate determinations, obtained by interpolation of concentration-inhibition curve.

II, some solvent partitioned fractions such as CH_2Cl_2 and EtOAc soluble fractions inhibited the oxidation of L-tyrosine catalyzed by tyrosinase with IC_{50} values of 183 and 67.0 $\mu\text{g/mL}$, respectively. However other fractions did not showed any activity at the highest concentration of 500 $\mu\text{g/mL}$. The EtOAc soluble fraction of the MeOH extract of *E. stolonifera* exhibited stronger inhibitory activity on the oxidation of L-tyrosine by the mushroom tyrosinase than that of arbutin, positive control with an IC_{50} value of 112 $\mu\text{g/mL}$. Column chromatography of the EtOAc fraction led us to the isolation of tyrosinase inhibitory phloroglucinol derivatives. These phloroglucinol derivatives were identified as phloroglucinol (1), eckstolonol (2), eckol (3), phlorofucofuroeckol A (4), and dieckol (5) on the basis of chemical and physicochemical evidences, and compared their spectral data with those in the literature as previously reported (Fukuyama *et al.*, 1985, 1989a, 1990; Nakamura *et al.*, 1996; Kang *et al.*, 2003). The chemical structures of the five phloroglucinol derivatives are presented in Fig. 1. IC_{50} values of compounds 1~5 on the oxidation of the L-tyrosine catalyzed by tyrosinase, were determined to be 92.8, 126, 33.2, 177, and 2.16 $\mu\text{g/mL}$, respectively, as shown at Table III. It was compared with kojic acid and arbutin, positive controls, with IC_{50} values of 6.32 and 112

Table II. Inhibitory activities of various fractions obtained from the MeOH extract of *E. stolonifera* on mushroom tyrosinase activity

Samples	IC_{50} ($\mu\text{g/mL}$) ^a
MeOH ex.	354.0
<i>n</i> -Hexane fr.	>500.0
CH_2Cl_2 fr.	183.0
EtOAc fr.	67.0
<i>n</i> -BuOH fr.	>500.0
H_2O fr.	>500.0
Arbutin	112.0
Kojic acid	6.32

^aInhibitory activity was expressed as the mean of 50% inhibitory concentrations of triplicate determinations, obtained by interpolation of concentration-inhibition curve.

Table III. Tyrosinase inhibitory activity of isolated compounds 1~5 from the EtOAc fraction of the MeOH extract of *E. stolonifera*

Compounds	IC_{50} ($\mu\text{g/mL}$) ^a
Phloroglucinol (1)	92.8
Eckstolonol (2)	126.0
Eckol (3)	33.2
Phlorofucofuroeckol A (4)	177.0
Dieckol (5)	2.16
Arbutin	112.0
Kojic acid	6.32

^aInhibitory activity was expressed as the mean of 50% inhibitory concentrations of triplicate determinations, obtained by interpolation of concentration-inhibition curve.




Fig. 1. The structures of the phlorotannins from *E. stolonifera*

$\mu\text{g/mL}$, respectively. All the compounds inhibited the monophenolase activity of the mushroom tyrosinase, and phloroglucinol (1) and eckol (3) showed stronger inhibitory activity than arbutin, and dieckol (5) displayed three times higher activity than kojic acid, a well-known tyrosinase inhibitor.

Lineweaver-Burk plots as shown in Fig. 2 and 3, depicted to determine the inhibitory modes of compounds 1~5, respectively. The lines, obtained from the uninhibited enzyme and from three different concentrations of the test compounds, intersected on the vertical axis (Fig. 2). The results demonstrate that compounds 1 and 2 exhibited competitive inhibition against oxidation of L-tyrosine catalyzed by mushroom tyrosinase. K_i values of phloroglucinol (1) and eckstolonol (2) were estimated to be 2.3×10^{-4} and 3.1×10^{-4} M, respectively. The lines, obtained with the uninhibited enzyme and from different concentrations of eckol (3), phlorofucofuroeckol A (4) or dieckol (5), intersected on the horizontal axis (Fig. 3). The results demonstrate that compounds 3, 4, and 5 exhibited non-competitive inhibitors for the oxidation of L-tyrosine catalyzed by mushroom tyrosinase. K_i values of 3, 4, and 5 which can be calculated from the slope or the $1/V$ -axis intercept, were 1.9×10^{-5} , 1.4×10^{-3} , and 1.5×10^{-5} M, respectively (Table IV).

A competitive inhibitor is a substance that combines with a free enzyme (E) in a manner that prevents substrate binding. That is, the inhibitor (I) and the substrate (S) are mutually exclusive, often because of true competition for the same site. A competitive inhibitor might be a nonmeta-




Fig. 2. Lineweaver-Burk plots for the inhibition of the mushroom tyrosinase by phloroglucinol (A) and eckstolonol (B). The reaction was done in the presence of phloroglucinol [final concentration of 0 (●), 125 (▼), 250 (■) and 500 μM (◆)] or eckstolonol [0 (●), 50 (◆), 100 (■) and 200 μM (▼)]. $V_{\text{max}} = 0.02 \Delta \text{OD}_{490}/\text{min}$, $K_m = 1.2 \times 10^{-3}$ M, $1/V : 1/(\Delta \text{OD}_{490}/\text{min})$.

Table IV. Tyrosinase inhibitory modes and inhibitory constants (K_i) of phloroglucinol derivatives isolated from *E. stolonifera*

Compounds	Mode of inhibition ^a	K_i value (M)
Phloroglucinol (1)	Competitive	2.3×10^{-4}
Eckstolonol (2)	Competitive	3.1×10^{-4}
Eckol (3)	Noncompetitive	1.9×10^{-5}
Phlorofucofuroeckol A (4)	Noncompetitive	1.4×10^{-3}
Dieckol (5)	Noncompetitive	1.5×10^{-5}

^aWith respect to the oxidation of L-tyrosine by mushroom tyrosinase.

bolizable analogue or derivative of the true substrate, or alternate substrate of the enzyme, or a product of the reaction. A classical noncompetitive inhibitor has no effect on substrate binding and *vice versa*. S and I bind




Fig. 3. Lineweaver-Burk plots for the inhibition of the mushroom tyrosinase by eckol (A), phlorofucofuroeckol A (B) and dieckol (C). The reaction was done in the presence of eckol [final concentration of 0 (●), 2.5 (◆), 5 (■) and 10 μM (▼)], phlorofucofuroeckol A [0 (●), 208 (▽), and 313 μM (▼)] or dieckol [0 (●), 5 (▼), and 10 μM (□)]. $V_{\text{max}} = 0.02 \Delta\text{OD}_{490}/\text{min}$, $K_m = 1.7 \times 10^{-3} \text{ M}$, $1/V : 1/(\Delta \text{OD}_{490}/\text{min})$.

reversibly, randomly, and independently at different sites (Segel, 1976).

Phloroglucinol derivatives, common secondary metabolite constituents of brown algae, are polymers of acetate-malonate derived 1,3,5-trihydroxybenzene (phloroglucinol)

(Higa, 1981). However, preincubation of the enzyme with these inhibitors in the absence of the substrate did not diminish enzyme activity significantly, suggesting that compounds 1-5 are inhibitors rather than inactivators of the enzyme (Khan and Andrawis, 1985).

Fukuyama *et al.* (1985; 1989a; 1989b; 1990) reported that phloroglucinol itself, and phlorethols did not show inhibitory activity on plasma plasmin at all, revealing the requirement of a dibenzo-1,4-dioxine ring for anti-plasmin inhibitory activity (Nakayama *et al.*, 1989; Hay, 1988). Phloroglucinol derivatives from *Eisenia bicyclis* showed a potent antioxidant effect which extended the induction time of autoxidation of methyl α -linolenate, seems to depend on the degree of polymerization of phloroglucinol. Generally, phloroglucinol derivatives of lower molecular weight were more effective than those of higher weight (Nakamura *et al.*, 1996). On the contrary, the feeding deterrent effect of marine herbivores (Taniguchi *et al.*, 1991; Hay, 1988; Altena and Steinberg, 1992; Boettcher and Targett, 1993; Targett *et al.*, 1995) and bactericidal activity (Nagayama *et al.*, 2002) of phloroglucinol derivatives tend to increase with polymerization of phloroglucinol. Although the mode of activity of phloroglucinol derivatives is still obscure; the interaction of phloroglucinol derivatives with bacterial proteins may play an important role in its bactericidal action (Stem *et al.*, 1996). Recently, Okada *et al.* reported the phloroglucinol derivatives from *E. bicyclis* exhibited inhibitory activity on glycation and α -amylase (Okada *et al.*, 2004). Although tyrosinase inhibitory activities of phloroglucinol derivatives such as phloroglucinol and eckol have been reported (Mitani, and Sakai, 1992; Onodera *et al.*, 1992), the tyrosinase inhibitors from *E. stolonifera* and their inhibitory kinetics have not been reported so far.

Nerya *et al.* (2003) suggested that one of the structural requirements for noncompetitive inhibitors on the tyrosinase is the presence of a resorcinol moiety, and that most of the competitive inhibitors have the ability to chelate copper in this enzyme, implying a possible inhibitory mechanism. All the phloroglucinol derivatives isolated from *E. stolonifera* contain resorcinol moieties, their tyrosinase inhibitory mode, however, exhibited two different types as described above. It will need the further study for the structure-activity relationships of tyrosinase inhibitors.

Dioxygen or reactive oxygen species (ROS) are required by the enzyme for the oxidation of L-tyrosine to 3-hydroxy-L-tyrosine (L-dopa) and then to L-dopaquinone (Schallreuter, 1999). Yokochi *et al.* (2003) reported that the vitamin B₆ inhibited the diphenolase activity by quenching ROS (probable singlet oxygen) generated during some reaction step of the diphenolase reaction. Further, they suggested that the inhibition is dependent on the quenching of ROS produced in some intermediary step of the tyrosinase

reaction. Previously, we investigated that the phloroglucinol derivatives from *E. stolonifera* inhibited total ROS generation (Kang *et al.*, 2004). Thus, it may be one of the probable inhibitory mechanisms that phloroglucinol derivatives inhibit tyrosinase by scavenging ROS produced during the catalytic cycle of the enzyme.

Safety is a primary consideration for tyrosinase inhibitors, especially for those used in food and cosmetic products, as these may be regularly utilized in unregulated quantities. In higher plants and fungi, tyrosinase inhibitors occur in various compounds, such as oxyresveratrol (Shin *et al.*, 1998), quercetin (Kubo and Kinst-Hori, 1999), (-) epigallocatechin 3-O-gallate (EGCG) (No *et al.*, 1999), and metallothionein (Goetghebeur and Kermasha, 1996), however, its inhibitors from edible alga *E. stolonifera* have not been found. Safety is a primary consideration for tyrosinase inhibitors, especially for those used in food and cosmetic products, as these may be regularly utilized in unregulated quantities. This work showed that phlorotannins, natural compounds found in an edible brown alga, could be involved in the control of pigmentation in plants and other organisms through inhibition of tyrosinase activity using L-tyrosine as a substrate.

ACKNOWLEDGEMENT

This research was supported by a grant (p-2004-02) from Marine Bioprocess Research Center of the Marine Bio 21 Center funded by the Ministry of Maritime Affairs & Fisheries, Republic of Korea.

REFERENCES

- Altena, A. and Steinberg, P. D., Are differences in the responses between North American and Australasian marine herbivores to phlorotannins due to differences in phlorotannin structure? *Biochem. Syst. Ecol.*, 20, 493-499 (1992).
- Andersen, S. O., Biochemistry of insect cuticle. *Annu. Rev. Entomol.*, 24, 29-61 (1979).
- Boettcher, A. A. and Targett, N. M., Role of polyphenolic molecular size in reduction of assimilation efficiency in *Xiphister mucosus*. *Ecology*, 74, 891-903 (1993).
- Choi, J. S., Lee, J. H., Park, H. J., Kim, H. G., Young, H. S., and Mun, S. I., Screening for antioxidant activity of plant and marine algae and its active principles from *Prunus davidiana*. *Kor. J. Pharmacogn.*, 24, 299-303 (1993).
- Fukuyama, Y., Miura, I., Kinzyo, Z., Mori, H., Kido, M., Nakayama, Y., Takahashi, M., and Ochi, M., Eckols, novel phlorotannins with a dibenzo-*p*-dioxin skeleton possessing inhibitory effects on α_2 -macroglobulin from the brown alga *Ecklonia kurome* OKAMURA. *Chem. Lett.*, 739-742 (1985).
- Fukuyama, Y., Kodama, M., Miura, I., Kinzyo, Z., Kido, M., Mori, H., Nakayama, Y., and Takahashi, M., Structure of an anti-plasmin inhibitor, eckol, isolated from the brown alga *Ecklonia kurome* OKAMURA and inhibitory activities of its derivatives on plasma plasmin inhibitors. *Chem. Pharm. Bull.*, 37, 349-353 (1989a).
- Fukuyama, Y., Kodama, M., Miura, I., Kinzyo, Z., Mori, H., Nakayama, Y., and Takahashi, M., Anti-plasmin inhibitor. V. Structures of novel dimeric eckols isolated from the brown alga *Ecklonia kurome* OKAMURA. *Chem. Pharm. Bull.*, 37, 2438-2440 (1989b).
- Fukuyama, Y., Kodama, M., Miura, I., Kinzyo, Z., Mori, H., Nakayama, Y., and Takahashi, M., Anti-plasmin inhibitor. VI. Structure of phlorofucofuroeckol A, a novel phlorotannin with both dibenzo-1,4-dioxin and dibenzofuran elements, from the brown alga *Ecklonia kurome* OKAMURA. *Chem. Pharm. Bull.*, 38, 133-135 (1990).
- Goetghebeur, M. and Kermasha, S., Inhibition of polyphenol oxidase by copper-metallothionein from *Aspergillus niger*. *Phytochemistry*, 42, 935-940 (1996).
- Han, E. S., Kim, J. W., Eom, M. O., Kang, I. H., Kang, H. J., Choi, J. S., Ha, K. W., and Oh, H. Y., Inhibitory effects of *Ecklonia stolonifera* on gene mutation on mouse lymphoma tk⁺ locus in L5178Y-3.7.2C cell and bone marrow micronuclei formation in ddY mice. *Environ. Mutagen. Carcinogen.*, 20, 104-111 (2000).
- Hay, M. E., Marine plant-herbivore interactions: the ecology of chemical defense. *Ann. Rev. Ecol. Syst.*, 19, 111-145 (1988).
- Higa, T., Phenolic substances. In Scheuer, P. J. (Ed.). *Marine Natural Products, Chemical and Biological Perspectives*. Vol. IV. Academic Press, New York, pp. 119-123 (1981).
- Kang, H. S., Chung, H. Y., Jung, J. H., Son, B. W., and Choi, J. S., A new phlorotannin from the brown alga *Ecklonia stolonifera*. *Chem. Pharm. Bull.*, 51, 1012-1014 (2003).
- Kang, H. S., Chung, H. Y., Kim, J. Y., Son, B. W., Jung, H. A., and Choi, J. S., Inhibitory phlorotannins from the edible brown alga *Ecklonia stolonifera* on total reactive oxygen species (ROS) generation. *Arch. Pharm. Res.*, 27, 194-198 (2004).
- Kim, Y. M., Yun, J., Lee, C. K., Lee, H. H., Min, K. R., and Kim, Y., Oxyresveratrol and hydroxystilbene compounds. Inhibitory effect on tyrosinase and mechanism of action. *J. Biol. Chem.*, 277, 16340-16344 (2002).
- Khan, V. and Andrawis, A., Inhibition of mushroom tyrosinase by tropolone. *Phytochemistry*, 24, 905-908 (1985).
- Kubo, I. and Kinst-Hori, I., Flavonols from saffron flower: tyrosinase inhibitory activity and inhibition mechanism. *J. Agric. Food Chem.*, 47, 4121-4125 (1999).
- Kurata, K., Taniguchi, K., Shiraiishi, K., Hayama, N., Tanaka, I., and Suzuki, M., Ecklonialactone-A and -B, two unusual metabolites from the brown algae *Ecklonia stolonifera* Okamura. *Chem. Lett.*, 267-270 (1989).
- Kurata, K., Taniguchi, K., Shiraiishi, K., and Suzuki, M., Ecklonialactones C-F from the brown alga *Ecklonia stolonifera*. *Phytochemistry*, 33, 155-159 (1993).
- Lee, J. H., Kim, N. D., Choi, J. S., Kim, Y. J., Heo, M. Y., Lim, S.

- Y., and Park, K. Y., Inhibitory effects of the methanolic extract of an edible brown alga, *Ecklonia stolonifera* and its component, phloroglucinol on aflatoxin B₁ mutagenicity *in vitro* (Ames test) and on benzo(a)pyrene or *N*-methyl *N*-nitrosourea clastogenicity *in vivo* (mouse micronucleus test). *Nat. Prod. Sci.*, 4, 105-114 (1998).
- Lee, J. H., Oh, H. Y., and Choi, J. S., Preventive effect of *Ecklonia stolonifera* on the frequency of benzo(a)pyrene-induced chromosomal aberrations. *J. Food Sci. Nutr.*, 1, 64-68 (1996).
- Lee, J. H., Park, J. C., and Choi, J. S., The antioxidant activity of *Ecklonia stolonifera*. *Arch. Pharm. Res.*, 19, 223-227 (1996).
- Mason, H. S., Structures and functions of the phenolase complex. *Nature*, 177, 79-81, (1956).
- Mayer, A. M., Polyphenol oxidase in plants - Recent progress. *Phytochemistry*, 26, 11-20 (1987).
- Mitani, Y. and Sakai, S., Eckols as tyrosinase inhibitors. *Jpn. Kokai Tokyo Koho*, JP 04235110 (1992).
- Nagayama, K., Iwamura, Y., Shibata, T., Hirayama, I., and Nakamura, T., Bactericidal activity of phlorotannins from the brown alga *Ecklonia kurome*. *J. Antimicro. Chemother.*, 50, 889-893 (2002).
- Nakamura, T., Nagayama, K., Uchida, K., and Tanaka, R., Antioxidant activity of phlorotannins isolated from the brown alga *Eisenia bicyclis*. *Fish. Sci.*, 62, 923-926 (1996).
- Nakayama, Y., Takahashi, M., Fukuyama, Y., and Kinzyo, Z., An anti-plasmin inhibitor, eckol, isolated from the brown alga *Ecklonia kurome* OKAMURA. *Agric. Biol. Chem.*, 53, 3025-3030 (1989).
- Nerya, O., Vaya, J., Musa, R., Izrael, S., Ben-Arie, R., and Tamir, S., Glabrene and isoliquiritigenin as tyrosinase inhibitors from Licorice Roots. *J. Agric. Food Chem.*, 51, 1201-1207 (2003).
- No, J. K., Soung, D. Y., Kim, Y. J., Shim, K. H., Jun, Y. S., Rhee, S. H., Yokozawa, T., and Chung, H. Y., Inhibition of tyrosinase by green tea components. *Life Sci.*, 65, PL 241-246 (1999).
- Okada, Y., Ishimaru, A., Suzuki, R., and Okuyama, T., A new phloroglucinol derivative from the brown alga *Eisenia bicyclis*: Potential for the effective treatment of diabetic complications. *J. Nat. Prod.*, 67, 103-105 (2004).
- Onodera, J., Yamamoto, T., Abe, K., and Ueno, N., Cosmetics containing phloroglucinol derivatives. *Jpn. Kokai Tokyo Koho*, Jp 06092835 (1994).
- Pérez-Bernal, A., Muñoz-Pérez, M. A., and Camacho, F., Management of facial hyperpigmentation. *Am. J. Clin. Derm.*, 1, 261-268 (2000).
- Schallreuter, K. U., A review of recent advances on the regulation of pigmentation in the human epidermis. *Cell Mol. Biol.*, 45, 943-949 (1999).
- Segel, I. H., Enzyme. In Segel, I. H. (ed.). *Biochemical Calculations*. John Wiley & Sons Inc., New York, pp. 246-256 (1976).
- Shin, N. H., Ryu, S. Y., Choi, E. J., Kang, S. H., Chang, I. M., Min, K. R., and Kim, Y., Oxyresveratrol as the potent inhibitor on dopa oxidase activity of mushroom tyrosinase. *Biochem. Biophys. Res. Commun.*, 243, 801-803 (1998).
- Stern, J., Hageman, A., Steinberg, P., and Magon, P., Phlorotannin-protein interactions. *J. Chem. Ecol.*, 22, 1877-1899 (1996).
- Taniguchi, K., Kurata, K., and Suzuki, M., Feeding-deterrent effect of phlorotannins from the brown alga *Ecklonia stolonifera* against the abalone *Haliotis discus* Hannai. *Nippon Suisan Gakkaishi*, 57, 2065-2071 (1991).
- Targett, N. M., Boettcher, A. A., Targett, T. E., and Vrolijk, N. H., Tropical marine herbivore assimilation of phenolic-rich plants. *Oecologia*, 103, 170-179 (1995).
- Whitaker, J. R., Polyphenol oxidase, In Wong, D. W. S. (ed.). *Food Enzymes, Structure and Mechanism*, Chapman & Hall, New York, pp. 271-307 (1995).
- Wilcox, D. E., Porras, A. G., Hwang, Y. T., Lerch, K., Winkler, M. E., and Solomon, E. I., Substrate analogue binding to the coupled binuclear copper active site in tyrosinase. *J. Am. Chem. Soc.*, 107, 4015-4027 (1985).
- Xu, Y., Stokes, A. H., Freeman, W. M., Kumer, S. C., Vogt, B. A., and Vrana, K. E., Tyrosinase mRNA is expressed in human substantia nigra. *Mol. Brain Res.*, 45, 159-162 (1997).
- Yokochi, N., Morita, T., and Yagi, T., Inhibition of diphenolase activity of tyrosinase by vitamin B₆ compounds. *J. Agric. Food Chem.*, 51, 2733-2736 (2003).
- Source : *Arch. Pharm. Res.*, (2004), 27(12): 1226-1232

