



저작자표시-비영리-변경금지 2.0 대한민국

이용자는 아래의 조건을 따르는 경우에 한하여 자유롭게

- 이 저작물을 복제, 배포, 전송, 전시, 공연 및 방송할 수 있습니다.

다음과 같은 조건을 따라야 합니다:



저작자표시. 귀하는 원저작자를 표시하여야 합니다.



비영리. 귀하는 이 저작물을 영리 목적으로 이용할 수 없습니다.



변경금지. 귀하는 이 저작물을 개작, 변형 또는 가공할 수 없습니다.

- 귀하는, 이 저작물의 재이용이나 배포의 경우, 이 저작물에 적용된 이용허락조건을 명확하게 나타내어야 합니다.
- 저작권자로부터 별도의 허가를 받으면 이러한 조건들은 적용되지 않습니다.

저작권법에 따른 이용자의 권리는 위의 내용에 의하여 영향을 받지 않습니다.

이것은 [이용허락규약\(Legal Code\)](#)을 이해하기 쉽게 요약한 것입니다.

[Disclaimer](#)

Thesis for the Degree of Master of Science

**Differential Daily Rhythms of Melatonin
in the Pineal Gland and Gut of Goldfish,
Carassius auratus in Response to Light**



Ji Yong Choi

Department of Marine Bioscience and Environment

The Graduate School

Korea Maritime and Ocean University

February 2016

**Differential Daily Rhythms of Melatonin
in the Pineal Gland and Gut of Goldfish,
Carassius auratus in Response to Light**

Advisor: Prof. Cheol Young Choi



A dissertation submitted in partial fulfillment of the requirements for
the degree of

Master of Science

Department of Marine Bioscience and Environment
Graduate School of Korea Maritime and Ocean University

February 2016

**Differential Daily Rhythms of Melatonin
in the Pineal Gland and Gut of Goldfish,
Carassius auratus in Response to Light**

A dissertation

by

Ji Yong Choi

Certified by

Cheol Young Choi

Thesis Advisor

Accepted by

Sung Hwoan Cho

Chairman

Young-Ung Choi

Member

Cheol Young Choi

Member

December 2015

CONTENTS

Contents	i
List of Tables	iii
List of Figures	iv
List of Abbreviations	vi
Abstract (in Korean)	vii
Abstract	1
1. Introduction	3
2. Materials and methods	6
2.1. Experimental fish, ophthalmectomy, and conditions	6
2.2. <i>In vitro</i> culture of pineal gland and whole gut cells	7
2.3. Quantitative PCR (QPCR)	9
2.4. Western blot analysis	12
2.5. Melatonin concentration level	12
2.6. H ₂ O ₂ concentration measurement	13
2.7. Statistical analysis	13
3. Results	14
3.1. Change of pineal gland AANAT2 expression levels	14
3.2. Change of whole gut AANAT2 expression levels	14
3.3. Change of pineal gland MT-R1 protein and mRNA expression levels	20
3.4. MT-R1 protein and mRNA expression levels	20
3.5. Melatonin concentration	25
3.6. AANAT2 and MT-R1 expression in response to thermal stress	28
3.7. H ₂ O ₂ concentration under thermal stress conditions	28

4. Discussion	35
Acknowledgements	39
References	40



List of Table

Table 1. Primers used for QPCR amplification 11



List of Figures

- Fig. 1.** Changes in the expression levels of AANAT2 mRNA (A) in the pineal gland, for 2 days using various photoperiods [12L:12L (Cont.), LL and DD] and ophthalmectomy (OP). Changes of expression levels of AANAT2 mRNA in the cultured pineal gland cells (B) using various photoperiods [12L:12L (Cont.), LL and DD] 16
- Fig. 2.** Changes in the expression levels of AANAT2 mRNA (A) in the whole gut, for 2 days using various photoperiods [12L:12L (Cont.), LL and DD] and ophthalmectomy (OP). Changes of expression levels of AANAT2 mRNA in the cultured whole gut cells (B) using various photoperiods [12L:12L (Cont.), LL and DD] 18
- Fig. 3.** Changes in the expression levels of MT-R1 protein (A) and MT-R1 mRNA (B) in the pineal gland, for 2 days using various photoperiods [12L:12L (Cont.), LL and DD] and ophthalmectomy (OP). Changes of expression levels of MT-R1 mRNA in the cultured pineal gland cells (C) using various photoperiods [12L:12L (Cont.), LL and DD] 21
- Fig. 4.** Changes in the expression levels of MT-R1 protein (A) and MT-R1 mRNA (B) in the whole gut, for 2 days using various photoperiods [12L:12L (Cont.), LL and DD] and ophthalmectomy (OP). Changes of expression

levels of MT-R1 mRNA in the cultured whole gut cells (C) using various photoperiods [12L:12L (Cont.), LL and DD]	23
Fig. 5. Changes in the levels of melatonin in the pineal gland (A), plasma (B) and whole gut (C) of goldfish under various photoperiods [12L:12L (Cont.), LL and DD] and ophthalmectomy (OP) as measured by plate reader	26
Fig. 6. Changes in the levels of AANAT2 mRNA in goldfish pineal gland (A) and in the whole gut (B) under different temperature conditions, 22°C (control) and 30 °C (thermal stress) under the natural photoperiod (SNP)	29
Fig. 7. Changes in the levels of MT-R1 mRNA in goldfish pineal gland (A) and in the whole gut (B) under different temperature conditions, 22°C (control) and 30°C (thermal stress) under thenatural photoperiod (12L:12D)	31
Fig. 8. Changes in the levels of H ₂ O ₂ in the pineal gland (A) and gut (B) of goldfish at 22°C (control) and 30°C (thermal stress), under natural photoperiod conditions (12L:12D)	33

List of Abbreviations

AANAT	aralkylamin N-acetyltransferase
ANOVA	analysis of variance
CAT	catalase
cAMP	cyclic AMP
Ct	threshold cycle
DD	continuous dark
ELISA	enzyme-linked immunosorbent assay
GAPDH	glyceraldehyde-3-phosphate dehydrogenase
GPX	glutathione peroxidase
H ₂ O ₂	hydrogen peroxidehydroxyl radicals
LD	12-h light:12-h dark
LL	continuous light
Melatonin	N-acetyl-5-methoxytryptamine
mRNA	messenger RNA
MT-R	melatonin receptor
OP	ophthalmectomy group
QPCR	quantitative real-time polymerase chain reaction
ROS	reactive oxygen species
RT	reverse transcription
RT-PCR	reverse transcription polymerase chain reaction
SCN	suprachiasmatic nucleus
SE	standard error
SOD	superoxide dismutase
TBS	tris-buffered saline
ZT	zeitgeber time

빛 조절에 의한 금붕어, *Carassius auratus* 송과체와 장 멜라토닌의 일주기 리듬 차이

최 지 용

한국해양대학교 대학원 해양생명환경학과

요 약

본 연구는 빛을 감지하여 일주성을 포함한 생체리듬 조절에 관여하는 호르몬인 멜라토닌(melatonin)이 빛을 감지하는 기관인 송과체에서 뿐만 아니라 빛의 영향을 직접적으로 받지 않는 기관인 장에서도 분비된다는 사실에 착안하여, 금붕어, *Carassius auratus*를 대상으로 송과체와 장에서 분비되는 멜라토닌의 일주기리듬 차이 여부를 확인하고자 수행되었다. 또한, 강력한 항산화 물질로 알려져 있는 멜라토닌이 분비기관(송과체와 장)에 따라 기능적인 측면에서 차이가 있는지에 대해서도 확인하고자 하였다.

1. 금붕어의 송과체와 장에서 각각 분비되는 멜라토닌의 일주기리듬 차이 여부를 확인하기 위하여 <멜라토닌 합성에 관여하는> aralkylamine N-acetyltransferase (AANAT)와 멜라토닌 수용체 유전자의 발현량 및 혈중 멜라토닌의 농도 변화를 관찰하였다. 금붕어의 송과체와 장에서

AANAT와 멜라토닌 수용체 cDNA를 각각 분리하였으며, 48 시간 동안 명·암기(LD), 명기(LL), 암기(DD) 및 안구적출(OP) 조건에서 금붕어의 송과체와 장 조직을 대상으로 이들 유전자의 발현량 변화를 분석하였다. LD 광 조건의 송과체에서는 AANAT와 멜라토닌 수용체 유전자가 낮 시간대(ZT8)에 비하여 밤 시간대(ZT20)에 높은 발현량을 나타내었고, 송과체 배양 실험(*in vitro*)에서도 비슷한 경향이 관찰되었다. DD 실험구는 LD 실험구에서의 AANAT와 멜라토닌 수용체 유전자의 발현 경향과 유사하게 나타났으나, AANAT와 멜라토닌 수용체 유전자의 발현량이 LD 실험구에 비하여 낮 시간대에 매우 높게 발현된 점으로 보아, 낮과 밤에 따른 일주기 경향이 약화된 것을 알 수 있었다. 특히, OP 실험구도 DD 실험구와 동일한 결과가 얻어졌는데, 이는 안구가 적출된 금붕어는 빛을 인지할 수 없기 때문에 DD 광 조건으로 인식하여 AANAT와 멜라토닌 수용체 유전자의 발현량이 DD 실험구에서의 결과와 동일한 것으로 사료된다.

한편, 장에서는 AANAT와 멜라토닌 수용체 유전자 발현량이 다른 모든 실험구에서 낮 시간대(ZT8)에 비하여 밤 시간대(ZT20)에 높게 나타났으며, 장 배양 실험(*in vitro*)에서도 비슷한 경향이 관찰되었다. 또한, 송과체, 혈장 및 장에서 멜라토닌의 일주기리듬을 분석한 결과, AANAT와 멜라토닌 수용체 유전자의 발현량 변화와 유사한 결과가 나타났다.

따라서 장에서의 AANAT와 멜라토닌 수용체 유전자 및 멜라토닌의 일주기리듬은 빛의 유무와는 상관없이 모든 실험구에서 동일한 일주기리듬을 보인 점으로 보아, 장에서 분비되는 멜라토닌은 빛에 의하여 일주기리듬이 조절되는 송과체와는 달리 빛에 의한 영향을 받지 않으며, 자체적인 멜라토닌의 생체리듬을 갖고 있을 것으로 판단된다.

2. 또한, 항산화 물질인 멜라토닌이 분비되는 기관(송과체와 장)에 따라 기능적인 차이가 존재하는지를 확인하기 위하여, 금붕어를 스트레스 환경

인 30℃의 고수온에 노출시킨 후, 송과체와 장에서 각각 분비되는 멜라토닌의 항산화 기능의 차이 여부를 조사하였다. 그 결과, 고수온(30℃)에 노출된 금붕어는 대조구(22℃)의 금붕어에 비하여 송과체와 장 조직 모두에서 AANAT와 멜라토닌 수용체 유전자의 발현량이 유의적으로 높게 나타났다. 산화스트레스로 인하여 체내에서 유도된 활성산소(H_2O_2) 농도 변화 또한 AANAT와 멜라토닌 수용체 유전자의 발현 경향과 유사하게 나타났으나, 송과체와 장 조직 간에 유의적인 차이는 없는 것으로 나타났다. 즉, 금붕어는 고수온 노출에 의하여 유도된 산화스트레스에 의하여 체내 활성산소(H_2O_2)가 증가하였고, 금붕어는 활성산소를 제거(감소)하는 동시에 산화스트레스 환경에 적응하기 위하여 강력한 항산화 물질인 멜라토닌을 송과체와 장에서 동시에 분비한 결과, AANAT와 멜라토닌 수용체 유전자의 발현량이 증가된 것으로 보인다.

따라서 고수온 환경에 노출됨으로써 유도된 산화스트레스를 제거하기 위한 멜라토닌의 항산화 기능은 분비되는 기관(송과체와 장)에 따라서는 차이가 없는 것으로 판단되었다.

**Differential Daily Rhythms of Melatonin
in the Pineal Gland and Gut of Goldfish,
Carassius auratus in Response to Light**

Ji Yong Choi


**Department of Marine Bioscience and Environment
Graduate School of Korea Maritime and Ocean University**

Abstract

The objectives of this study were to test the effects of light on melatonin rhythms in the pineal gland and gut of goldfish (*Carassius auratus*) and to investigate whether melatonin function differed in these two tissues, which are respectively photosensitive and non-photosensitive. Rhythms were evaluated by measuring aralkylamine N-acetyltransferase (*AANAT2*) and melatonin receptor 1 (MT-R1) mRNA expression and melatonin concentration in the pineal gland, gut (*in vivo*), and cell cultures of the two tissues (*in vitro*). Compared to control, pineal gland melatonin secretion was higher at night, whereas the 24-h dark and ophthalmectomy groups maintained higher *AANAT2* and MT-R1 mRNA expression during the day. Melatonin levels and *AANAT2* and MT-R1 mRNA expression in the gut were also the

highest at night, but the 24-h light, dark, and ophthalmectomy groups did not significantly differ from control. Furthermore, we measured *AANAT2* and MT-R1 mRNA expression in high temperature water (30 °C) to investigate differences in the antioxidant capacity of pineal gland versus gut melatonin. Melatonin and H₂O₂ levels, as well as *AANAT2* and MT-R1 mRNA expression, were all higher in the two tissues under thermal stress, compared with their levels at 22 °C. Taken together, our results suggest that light has no effect on melatonin patterns in the gut, which appears to exhibit its own circadian rhythm, but both gut and pineal gland melatonin exhibit similar antioxidant function.

Keywords: *AANAT2*; circadian rhythms; gut; melatonin; ophthalmectomy; pineal gland



1. Introduction

Light is one of the most important environmental factors to influence nearly all organisms (Migaud et al., 2006). Generally, in vertebrates, including fish, photoreceptors in the retina process external light signals and transmits the information to the hypothalamic suprachiasmatic nucleus (SCN). From the SCN, signals are then transferred to photosensitive organs (Yoshikawa and Oishi, 1998; Yamazaki et al., 1999).

A major light-dependent hormone is melatonin (N-acetyl-5-methoxytryptamine), which plays a major role in controlling fish biorhythms as a neuroendocrine communicator (Falcón et al., 2007). Besides its function as a regulator of circadian rhythms, melatonin also performs various physiological functions, such as removing free radicals, enhancing the immune system, and reducing oxidative stress in fish (Wu and Swaab, 2005; Besseau et al., 2006; Kim et al., 2014). Melatonin acts by binding to melatonin receptors (MT-Rs), belonging to the G-protein coupled receptor superfamily and widely distributed in the vertebrate diencephalon and central nervous system (Reppert et al., 1996; Park et al., 2013). Among the MT-Rs, MT-R1 is implicated in the seasonal changes to vertebrate physiology and behavior, including those of fish, and directly affects circadian rhythms (Reppert et al., 1996).

Melatonin is secreted mainly at night by the pineal gland and retina, and regulated by arylalkylamine N-acetyltransferase (AANAT). AANAT is well known as a rate-limiting enzyme of the melatonin biosynthetic pathway (Klein et al., 2002). During night time, depolarization of light receptors results in an influx of Ca^{2+} and cyclic AMP (cAMP) into the cell, which then phosphorylates AANAT, increasing its intracellular amount and activity (Iuvone et al., 2005; Klein, 2007).

Although melatonin is widely accepted as a major regulator of vertebrate

biorhythms, the hormone has been also found in several non-photosensitive tissues (Bubenik, 1980; Vakkuri et al., 1985; Bubenik and Pang, 1997). One notable example is the gut, which was reported to secrete more melatonin than the pineal gland (Bubenik, 2002). In addition, melatonin secretion and AANAT expression patterns in the gut appear to vary depending on the species (Velarde et al., 2010). Previous studies have shown that the melatonin generating system in the gut and pineal gland of the major carp *Catla catla* may not be under photo-thermal regulation due to daily and seasonal photoperiod changes (Mukherjee et al., 2014a, b). Despite these recent insights, the molecular endocrinological mechanism underlying the autonomous circadian rhythm of gut melatonin regulated by light remains poorly understood.

Previous studies have shown that melatonin acts as a direct scavenger of free radicals. Reiter et al. (2000) suggested that melatonin increases the expression and activity of antioxidants, such as superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPX), and removes internally generated reactive oxygen species (ROS), including superoxide (O_2^-), hydrogen peroxide (H_2O_2), hydroxyl radicals (OH^-), and singlet oxygen (1O_2). However, little is known about the role of melatonin in photosensitive and non-photosensitive tissues, such as the pineal gland and gut, under thermal and salinity stress conditions.

Goldfish *Carassius auratus* is a popular ornamental fresh water fish and an ideal model to study the mechanisms of circadian rhythms and photo-response in any melatonin synthesizing tissue, because it maintains close contact with environmental light, due to its natural surface dwelling habit. To study the differences in the levels of AANAT2 and MT-R1 mRNA and melatonin concentrations between the photosensitive pineal tissue and non-photosensitive gut tissue, the goldfishes were reared under different

photoperiod conditions. In addition, this study also analyzed how the absence of the retina affected the circadian rhythms of pineal gland and gut melatonin after ophthalmectomy (OP). Finally, to ascertain whether gut melatonin shared antioxidant function as pineal gland melatonin, this study compared and analyzed the expression patterns of pineal gland and gut AANAT2 and MT-R1 mRNA, as well as the levels of H₂O₂ under thermal stress conditions (temperature increased from 22°C to 30°C in daily increments of 2°C).



2. Materials and methods

2.1. Experimental fish, ophthalmectomy, and conditions

For each experiment, common goldfish ($n = 720$, mean length, 8.1 ± 0.4 cm; mean mass, 16.5 ± 0.6 g) were purchased from a commercial aquarium (Choryang, Busan, Korea) and were allowed to acclimate for 2 weeks in eleven 300-L circulation filter tanks in the laboratory. Five tanks with 60 individuals each were used for light response and thermal stress experiments, and one tank with 60 individuals for cell culture experiment. Both experiments were duplicated.

Experimental groups [control (12L:12D), continuous light (LL), continuous dark (DD), ophthalmectomy (OP), and thermal stress] were exposed to light using a white fluorescent bulb (27 W, wavelength range 350–650 nm); placed 40 cm above the water surface. The light intensity at the water surface was approximately 0.96 W/m^2 . Goldfish were reared with automatic temperature regulation systems (JS-WBP-170RP; JohnsamCo., Seoul, Korea) and allowed to acclimatize to the experimental conditions for 24 h. Water temperature was maintained at 22°C . All the fish were provided commercial feed daily at 0900 and 1700.

For the light response study, the groups were used such as a control group (12L:12D) and three experimental groups (LL, DD, and OP). The control group was exposed to a 12-h light:12-h dark cycle (12L:12D) (lights were turned on at 0700 and turned off at 1900). The LL experimental group was exposed to 24-h light; the DD experimental group was exposed to 24-h dark; and the OP experimental group was exposed to 12L:12D.

Furthermore, the 120 fish in the OP experiment group were subjected to ophthalmectomy under anesthesia with 200 mg/L tricaine methane sulfonate (MS-222; Sigma-Aldrich, St. Louis, MO, USA). An incision was made around the eye, which was then elevated using a forceps, and the optic nerve was cut to remove the eye completely as described by Martinez-Chavez and Migaud (2009). These fish were placed in 300-L circulation filter tanks under 12L:12D period at 22°C for one week. No mortalities were observed.

In the thermal stress experiment group, fish were reared with a 12L:12D photoperiod under a white fluorescent bulb, with lights on at 0700 and lights off 1900. Fishes were all initially exposed to 22°C water; this temperature was maintained for the control group, but increased from 22°C to 30°C in daily increments of 2°C for the experimental groups.

The fish were killed by spinal transection (first sampling at 1100; ZT4) at 4-h sampling intervals (ZT4, ZT8, ZT12, ZT16, ZT20, and ZT24) to collect the pineal gland, whole gut, and blood under dim light using an attenuated white fluorescent bulb. The fish were anesthetized with 200 mg/L tricaine methane sulfonate (MS-222; Sigma-Aldrich, St. Louis, MO, USA) to minimize stress prior to blood collection. Blood was collected rapidly from the caudal vein using a 1-mL syringe coated with heparin. Plasma samples were separated with centrifugation (4°C, 1,000 ×g, 15 min) and stored at -80°C until analysis. The pineal gland and the gut were removed from the fish, immediately frozen in liquid nitrogen, and stored at -80°C until analysis.

2.2. *In vitro* culture of pineal gland and whole gut cells

For cultures, pineal gland and whole gut tissue were quickly removed and

placed in 3 mL dispersion buffer that was kept on ice for 1 min (dispersion buffer: pH 7.4, Dulbecco's phosphate-buffered saline, without calcium chloride and magnesium chloride, containing 100 U/mL penicillin, 100 µg/mL streptomycin and 2.5 µg/mL fungizone; GIBCOBRL, Rockville, MD, USA). The isolated pineal gland and whole gut tissues were then transferred to 6 mL of fresh dispersion buffer containing 0.25% trypsin (Type II-S from porcine pancreas; Sigma, USA). Next, tissue samples were cut into small pieces using a single-edged razor after connective tissues and other impurities were removed. The minced tissues of the two organs were transferred to a flask and incubated for 10 min at room temperature with slow stirring. The pineal gland and whole gut tissue mixture was filtered, and the culture medium was added (neuro basal medium, without L-glutamine, containing 100 U/mL penicillin, 100 µg/mL streptomycin, 2.5 µg/mL fungizone and 1% fetal bovine serum, Gibco-BRL; to adjust the medium osmolarity to goldfish plasma osmolarity, 353 mOs). The cell suspension was centrifuged at 800 ×g for 10 min, and the cells were then resuspended in fresh culture medium. Cells from the pineal gland and gut of 60 goldfish were counted and adjusted to a final concentration of approximately 1.93×10^6 live pineal gland cells/mL and 4.51×10^6 live gut cells/mL (as determined by trypan blue exclusion) in supplemented media.

The pineal gland and gut cells at a concentration of approximately 1.2×10^6 cells/mL were added in 24-well culture plates (SPL Life Sciences, Seoul, South Korea) and incubated at 22°C, 100% humidity, and 5% atmospheric CO₂ in a culture chamber (HB-302L; Hanbaek Scientific Co., Gyeonggi, Korea). Incubation started at 0700, and samples were collected at ZT4, ZT8,

ZT12, ZT16, ZT20, and ZT24. Thus, the first collection time was at the first ZT4 (1100) to second ZT24 (0700). During the experimental period (2 days), the pineal gland and gut cells were exposed to 12L:12D (control group), LL, and DD using a dim white fluorescent bulb placed 50 cm above the cell culture plate. After the end of incubation, a cell dissociation reagent (trypsin/EDTA) was used to degrade the adhesion between the cells and wells. Then, cell suspension was centrifuged at 800 ×g for 10 min, and the supernatant was removed and stored at −80°C until total RNA extraction and analysis.

2.3. Quantitative PCR (QPCR)

To carry out this study, the experiments has considered the recommendations of the MIQE (Minimum Information for Publication of Quantitative Real-time PCR Experiments) (Bustin et al., 2009). Total RNA was extracted from the pineal gland and whole gut using a TRIzol kit (Gibco/BRL, USA) according to manufacturer protocol. To obtain cDNA for QPCR, reverse transcription was performed using M-MLV reverse transcriptase (Promega, USA), again following manufacturer protocol. QPCR was conducted to determine the relative expression levels of MT-R1 (GenBank accession no. [AB481372](#)), AANAT2 (GenBank accession no. [GU205782](#)), β-actin (GenBank accession no. [AB039726](#)), and GAPDH (GenBank accession no. [AY641443](#)) mRNA in the pineal gland and whole gut. The primers used for QPCR are shown in Table 1. These primers were designed for each gene using the Beacon Designer software (Bio-Rad, Hercules, CA, USA). Primer alignment was performed with the BLAST

database to ensure primer specificity. PCR was conducted using a Bio-Rad CFX96™ Real-time PCR Detection System (Bio-Rad, Hercules, USA) and iQ™ SYBR Green Supermix (Bio-Rad, Hercules, USA), following manufacturer's protocol. The thermo cycling profile was as follows: 95°C for 5 min, followed by 50 cycles each of 95°C for 20 s and 55°C for 20 s. Experiments were duplicated with β -actin and GAPDH, and all data were expressed relative to the calculated β -actin and GAPDH threshold cycle (Δ Ct) levels.



Table 1. Primers used for QPCR amplification

Genes (accession no.)	Primer	DNA sequences
MT-R1 (AB481372)	Foeward	5'-GGT TGG CAG TAG CGA TTT-3'
	Reverse	5'-GGT TGG CAG TAG CGA TTT-3'
<i>AANAT2</i> (GU205782)	Foeward	5'-GAA TGT CCA CTC ACC CTT G-3'
	Reverse	5'-TCT CAT CGC TTC CTG TTC-3'
β -actin (AB039726)	Foeward	5'-TTC CAG CCA TCC TTC CTA-3'
	Reverse	5'-TAC CTC CAG ACA GCA CAG-3'
GAPDH (AY641443)	Foeward	5'-GGC CGC TAC AAG GGA GAT GT-3'
	Reverse	5'-CCG GTA GAC TCG ACT ACA TA -3'



2.4. Western blot analysis

Total protein isolated from the pineal gland and whole gut of goldfish was extracted using a T-PER[®] Tissue Protein Extraction Reagent (Thermo Fisher Scientific, Inc., USA) following manufacturer protocol. Twenty-five μ g total protein was loaded per lane onto Mini-PROTEAN[®] TGX[™] Gels (Bio-Rad, USA). For reference, a protein ladder (Bio-Rad, USA) was also used. Samples were electrophoresed at 180V. The gels were immediately transferred to a 0.2 μ m polyvinylidene difluoride (PVDF) membrane (Bio-Rad, USA), set at 85 V for 3 min, using the Trans-Blot[®] Turbo[™] Transfer System. There after, the membranes were blocked with 5% skim milk in 0.04% Tris-buffered saline with Tween (TTBS) for 45 min and subsequently washed in TTBS. The membranes were incubated with MT1 antibodies [Novusbiochemicals[®] Melatonin Receptor 1A Antibody (NBP1-28912); dilution, 1:4,000; approximately 45 kDa] for 2 hours, followed by horseradish peroxidase-conjugated anti-rabbit IgG secondary antibodies (dilution, 1:4,000; Bio-Rad, USA) for 60 min. As an internal control, incubation was performed with β -tubulin primary antibody (dilution, 1:4,000; ab6046, Abcam, UK), followed by horseradish peroxidase-conjugated anti-rabbit IgG secondary antibodies (dilution, 1:4,000; Bio-Rad) for 60 min. Bands were detected using a sensitive electrochemiluminescence (ECL) system (ECL Advance; GE Healthcare Life Sciences, Uppsala, Sweden) and exposed for 2 min using a Molecular Imager[®] ChemiDoc[™] XRS+ System (Bio-Rad, USA).

2.5. Melatonin concentration level

Pineal gland and whole gut samples were homogenized and centrifuged (4°C, 1,500 \times g, 15 min) and plasma samples were separated by centrifugation

(4°C, 1,000 ×g, 15 min). The melatonin levels of tissue and plasma were analyzed using immunoassays using Fish Melatonin ELISA kit (Catalog no. MBS013211; Mybiosource, San Diego, CA, USA). An anti-antibody that was specific to the melatonin antibody was pre-coated onto a microplate. Next, 100 µL tissue supernatant, 50 µL HRP-conjugate, and 50 µL antibody were added to each well. These were mixed and then incubated for 2 h at 37°C. Following the last wash, any remaining Wash Buffer was aspirated or decanted off, and 50 µL each of substrates A and B were added to each well. These substrate solutions were then incubated for 15 min at 37°C in the dark, during which they changed from colorless to dark blue. Following incubation, 50 µL stop solution was added to each well, changing the color from blue to yellow. The optical density of the solution in each well was then determined within 10 min, using a microplate reader set to 450 nm.

2.6. H₂O₂ concentration measurement

The H₂O₂ concentrations were measured using Peroxide Detect kit (Sigma-Aldrich, St.Louis, MO). Absorbance was measured at 560 nm, and the concentration of H₂O₂ was interpolated from a standard curve. The concentrations were expressed as nmol H₂O₂/g of tissue.

2.7. Statistical analysis

All data were analyzed using the SPSS statistical package (version 10.0; SPSS Inc., USA). One-way ANOVA followed by Tukey's post-hoc test was used to compare differences in the data ($p < 0.05$). The values are expressed as the means ± standard error (SE).

3. Results

3.1. Change of pineal gland AANAT2 expression levels

In all of the experimental groups (*in vivo*), AANAT2 mRNA expression was generally higher at night than during the day and was significantly higher at ZT20 (Fig 1). Within the DD and OP groups, AANAT2 mRNA expression significantly decreased during the second day cycle, compared with the first day cycle. When comparing across experimental and control groups, the present study were founded that both the DD and OP groups exhibited higher AANAT2 mRNA expression than the control group. Finally, the LL group exhibited significantly lower AANAT2 mRNA expression than the control as well as the DD and OP groups, while the latter two did not significantly differ from each other.

The QPCR results using *in vitro* mRNA (extracted from pineal gland cell culture) generally accorded with the *in vivo* experiments. Again across all experimental groups, AANAT2 mRNA expression was higher at night than during the day. In the DD group, AANAT2 mRNA expression also significantly decreased in the second day cycle compared with the first day cycle. Lastly, compared with control, AANAT2 mRNA expression was significantly higher in the DD group and lower in the LL group.


3.2. Change of whole gut AANAT2 expression levels

The results of the QPCR analysis on *in vivo* gut mRNA exhibited both similarities to, and differences from, mRNA patterns in the pineal gland (Fig. 2). First, AANAT2 mRNA expression was also higher at night than during

the day across all experimental groups, with the ZT20 time points exhibiting significantly higher expression levels; *in vitro* results exhibited the same pattern. However, in contrast to results from the pineal gland, AANAT2 mRNA expression did not differ between the first and second day cycles for both the DD and the OP groups. Furthermore, AANAT2 mRNA expression patterns did not significantly differ across any of the experimental groups, which was also the case with the *in vitro* results.



(A) AANAT2 mRNA expression in whole gut (*in vivo*)



(B) AANAT2 mRNA expression in whole gut (*in vitro*)






Fig. 1. Changes in the expression levels of AANAT2 mRNA (A) in the pineal gland, for 2 days using various photoperiods [12L:12L (Cont.), LL and DD] and ophthalmectomy (OP). Changes of expression levels of AANAT2 mRNA in the cultured pineal gland cells (B) using various photoperiods [12L:12L (Cont.), LL and DD]. The expression level of each sample was normalized with respect to the β -actin and GAPDH signal and is expressed as the relative expression level. The white bar represents the photophase and the black bar represents the scotophase. Values with different characters are same circadian time in fish exposed to the different experimental groups ($p < 0.05$). Numbers indicate significant differences between and within the experimental groups ($p < 0.05$). Data are expressed as mean \pm standard error (SE; $n = 5$).



(A) AANAT2 mRNA expression in whole gut (*in vivo*)



(B) AANAT2 mRNA expression in whole gut (*in vitro*)




Fig. 2. Changes in the expression levels of AANAT2 mRNA (A) in the whole gut, for 2 days using various photoperiods [12L:12L (Cont.), LL and DD] and ophthalmectomy (OP). Changes of expression levels of AANAT2 mRNA in the cultured whole gut cells (B) using various photoperiods [12L:12L (Cont.), LL and DD]. The expression level of each sample was normalized with respect to the β -actin and GAPDH signal and is expressed as the relative expression level. The white bar represents the photophase and the black bar represents the scotophase. Values with different characters are same circadian time in fish exposed to the different experimental groups ($p < 0.05$). Numbers indicate significant differences between and within the experimental groups ($p < 0.05$). Data are expressed as mean \pm standard error (SE; $n = 5$).



3.3. Change of pineal gland MT-R1 protein and mRNA expression levels

MT-R1 protein and mRNA expression (extracted from the pineal gland *in vivo*) were higher at night than during the day in all experimental groups, and was significantly higher at ZT20. In the DD and OP groups, MT-R1 mRNA expression also significantly decreased in the second day cycle compared with the first-day cycle. MT-R1 mRNA expression measured from *in vitro* pineal gland cells exhibited the same patterns of being higher at night and lower during the second day cycle (Fig. 3).


Compared to control, the DD and OP groups exhibited higher MT-R1 protein and mRNA expression, while the LL group exhibited significantly lower expression. Expression levels in the LL group were also significantly lower than the DD and OP groups, which did not significantly differ from each other. The results of *in vitro* experiments again mirrored these expression patterns, with DD groups expressing significantly higher levels of MT-R1 mRNA and LL groups expressing significantly lower levels.

3.4. MT-R1 protein and mRNA expression levels

MT-R1 protein and mRNA expression (extracted from goldfish gut *in vivo*) were higher at night than during the day time in all experimental groups, and were significantly higher at ZT20, a pattern that was mirrored by the results of the *in vitro* experiments. In contrast to the pineal gland, MT-R1 mRNA expression in the DD and OP groups did not significantly differ between the first day cycle and the second day cycle. Moreover, MT-R1 protein and mRNA expressions exhibited no significant difference across all experimental groups. The *in vitro* experiments also found no difference across the experimental groups (Fig. 4).


(A)

Western blot (pineal gland)




(B)

MT-R1 mRNA expression in pineal gland (*in vivo*)



(C)

MT-R1 mRNA expression in pineal gland (*in vitro*)




Zeitgeber time (h)

Fig. 3. Changes in the expression levels of MT-R1 protein (A) and MT-R1 mRNA (B) in the pineal gland, for 2 days using various photoperiods [12L:12L (Cont.), LL and DD] and ophthalmectomy (OP). Changes of expression levels of MT-R1 mRNA in the cultured pineal gland cells (C) using various photoperiods [12L:12L (Cont.), LL and DD]. The expression level of each sample was normalized with respect to the β -actin and GAPDH signal and is expressed as the relative expression level. The white bar represents the photophase and the black bar represents the scotophase. Values with different characters are same circadian time in fish exposed to the different experimental groups ($p < 0.05$). Numbers indicate significant differences between and within the experimental groups ($p < 0.05$). Data are expressed as mean \pm standard error (SE; $n = 5$).




(A)

Western blot (whole gut)




(B)

MT-R1 mRNA expression in whole gut (*in vivo*)



(C)

MT-R1 mRNA expression in whole gut (*in vitro*)



Zeitgeber time (h)

Fig. 4. Changes in the expression levels of MT-R1 protein (A) and MT-R1 mRNA (B) in the whole gut, for 2 days using various photoperiods [12L:12L (Cont.), LL and DD] and ophthalmectomy (OP). Changes of expression levels of MT-R1 mRNA in the cultured whole gut cells (C) using various photoperiods [12L:12L (Cont.), LL and DD]. The expression level of each sample was normalized with respect to the β -actin and GAPDH signal and is expressed as the relative expression level. The white bar represents the photophase and the black bar represents the scotophase. Values with different characters are same circadian time in fish exposed to the different experimental groups ($p < 0.05$). Numbers indicate significant differences between and within the experimental groups ($p < 0.05$). Data are expressed as mean \pm standard error (SE; $n = 5$).




3.5. Melatonin concentration

Melatonin concentrations in all tested samples (pineal gland, plasma, and gut) were significantly higher concentration at night than during the day (Fig. 5). The pineal gland and plasma exhibited similar patterns, with melatonin concentrations generally lower in the LL group than in the DD and OP groups. Additionally, melatonin concentrations in the latter two groups also significantly decreased in the second day cycle compared with the first day cycle. In contrast, melatonin concentrations in the gut did not significantly differ between the first day and the second day cycle.




(A)

Pineal gland melatonin




(B)

Plasma melatonin



(C)

Gut melatonin



Zeitgeber time (h)

Fig. 5. Changes in the levels of melatonin in the pineal gland (A), plasma (B) and whole gut (C) of goldfish under various photoperiods [12L:12L (Cont.), LL and DD] and ophthalmectomy (OP) as measured by plate reader. The white bar represents the photophase and the black bar represents the scotophase. Values with different characters are same circadian time in fish exposed to the different experimental groups ($p < 0.05$). Numbers indicate significant differences between and within the experimental groups ($p < 0.05$). Data are expressed as mean \pm standard error (SE; $n = 5$).



3.6. AANAT2 and MT-R1 expression in response to thermal stress

AANAT2 and MT-R1 mRNA expression in both the pineal gland and gut of the thermal stress group (30°C) were significantly higher than expression in the control group (22°C). For both groups, AANAT2 and MT-R1 mRNA expression were higher at night than during the day, and were significantly higher at ZT20 (Figs. 6 and 7).


3.7. H₂O₂ concentration under thermal stress conditions

The H₂O₂ concentrations in the pineal gland and gut of the thermal-stress group (30°C) were significantly higher than those of the control group (22°C) (Fig. 8). The concentrations in the pineal gland and gut did not differ significantly between the first day and second day cycles.




(A)

AANAT2 mRNA expression in pineal gland



(B)

AANAT2 mRNA expression in whole gut




Zeitgeber time (h)

Fig. 6. Changes in the levels of AANAT2 mRNA in goldfish pineal gland (A) and in the whole gut (B) under different temperature conditions, 22°C (control) and 30 °C (thermal stress) under the natural photoperiod (SNP). The expression level of each sample was normalized with respect to the β -actin and GAPDH signal and is expressed as the relative expression level. Values with different characters are at same circadian time ($p < 0.05$). Numbers indicate significant differences between and within the experimental groups ($p < 0.05$). Data are expressed as mean \pm standard error (SE; $n = 5$).



(A)

MT-R1 mRNA expression in pineal gland



(B)

MT-R1 mRNA expression in whole gut





Fig. 7. Changes in the levels of MT-R1 mRNA in goldfish pineal gland (A) and in the whole gut (B) under different temperature conditions, 22°C (control) and 30°C (thermal stress) under the natural photoperiod (12L:12D). The expression level of each sample was normalized with respect to the β -actin and GAPDH signal and is expressed as the relative expression level. Values with different characters are at same circadian time ($p < 0.05$). Numbers indicate significant differences between and within the experimental groups ($p < 0.05$). Data are expressed as mean \pm standard error (SE; $n = 5$).



(A)

H₂O₂ concentration in pineal gland



(B)

H₂O₂ concentration in whole gut




Fig. 8. Changes in the levels of H₂O₂ in the pineal gland (A) and gut (B) of goldfish at 22°C (control) and 30°C (thermal stress), under natural photoperiod conditions (12L:12D). Values with different characters are at the same circadian time ($p < 0.05$). Numbers indicate significant differences between and within the experimental groups ($p < 0.05$). Data are expressed as mean \pm standard error (SE; $n = 5$).



4. Discussion

In this study, it found that pineal gland and gut melatonin shared the same functions as a circadian rhythms regulator and an antioxidant, despite differences in the photosensitivity of the originating tissue. The presence or absence of light, however, did have different effects on melatonin, depending on whether it was secreted by the pineal gland (photosensitive) or the gut (non-photosensitive).

First, both *in vivo* and *in vitro* experiments confirmed that AANAT2 and MT-R1 mRNA expression were significantly higher at night (ZT20) in the pineal gland and gut, across all experimental groups. In the pineal gland, these patterns were significantly affected by the presence or absence of light, while patterns of expression in the gut remained unaffected by light. Similar to the results of this study, Coon et al. (1999) found that AANAT2 mRNA expression significantly decreased in the pineal gland of pike *Esox lucius* exposed to continuous light for 24 h. Moreover, Gothilf et al. (1999) also found that AANAT2 mRNA expression significantly increased in the pineal gland of zebrafish *Danio rerio* exposed to continuous dark for 24 h. Because continuous dark conditions resulted in increased AANAT2 and MT-R1 mRNA expression, this suggests that the constant dark environment controls melatonin synthesis through the promotion of AANAT2 mRNA expression, resulting in higher levels of MT-R1 expression as well. Although less well studied, this results on gut melatonin secretion patterns also have some corroboration in previous research: Vakkuri et al. (1985) reported that the secretion of gut melatonin in the pigeon was unaffected by exposure to light at midnight vs. exposure to a natural photoperiod.

In this study, goldfish subjected to ophthalmectomy had similar patterns of gut AANAT2 and MT-R1 mRNA expression compared with those exposed to

continuous light or dark. The gut melatonin possesses a circadian rhythm unaffected by light, and therefore unaffected by the presence or absence of the eye (retina). This seems logical given that the gut cannot directly receive light signals from the retina, whereas the more closely situated pineal gland is known to do so. In other words, this study suggests that the gut can regulate its own circadian rhythm, without using light signals, through the expression of AANAT2 and MT-1R mRNA. Further supporting this conclusion, the oscillation amplitude of gut AANAT2 and MT-R1 mRNA was maintained without reference to light, or the first and second day circadian cycles.

In contrast, pineal gland and plasma AANAT2 and MT-R1 mRNA were significantly lower during the second day cycle compared with the first day cycle (ZT4–ZT24). This decrease over time is not light sensitive, as it occurred in all of the experimental groups (LL, DD, and OP). A similar decrease in the oscillation amplitude of AANAT2 mRNA expression and activity was found in chickens exposed to continuous light or dark conditions (Klein et al., 1997). This pattern occurred here because the goldfish gradually adapted to either continuous light or continuous dark.

This study also determined that the daily secretion pattern of melatonin in the pineal gland, plasma, and gut were similar to the observed changes in AANAT2 and MT-R1 mRNA expression across all experimental groups. In particular, these results found no difference in melatonin concentrations between the DD and OP groups, as fish in the OP condition was effectively deprived of light signals, similar to the DD condition. Additionally, melatonin daily rhythms in the pineal gland and plasma are similar to each other, but gut melatonin rhythms observed constantly across all experimental groups (LL, DD, or OP).

Findings on gut melatonin are supported by a previous study in rats,

which also observed a constant daily rhythm of gut melatonin, even after a pinealectomy (Bubenik & Brown, 1997). Again, these results suggest that the gut possesses its own biorhythms for melatonin secretion and the synthesis of melatonin and AANAT2 in the gut is governed by a circadian clock and entrained by non-photic cues.

Despite differences in light sensitivity, this study found that gut and pineal gland melatonin both exhibited antioxidant function. In both organs, AANAT2 and MT-R1 mRNA expression increased in response to thermal stress, suggesting that the oxidative stress induced by high water temperatures caused an antioxidant response in the fish. Several studies corroborate our findings. For example, Choi et al. (2014) also found that MT-R1 mRNA expression significantly increased in goldfish as an antioxidant response to increased water temperatures. In addition, a study on trout found that high water temperature appears to regulate AANAT2 activity, thereby increasing melatonin synthesis (Benyassi et al., 2000).

In order to study the effects of oxidative stress on goldfish under thermal stress conditions, this experiment measured the levels of H₂O₂ in the pineal gland and gut. Significantly higher H₂O₂ levels were noted in the tissues in the thermal stress groups (30°C) than in tissues in the control groups (22°C). These results were in agreement with those reported by Nakano et al. (2014), in which the oxidative stress on coho salmon *Oncorhynchus kisutch* was higher under high-temperature conditions, and as a result, the levels of plasma lipid peroxides were also high. Our results suggested that the pineal gland and gut exhibit antioxidant functions under thermal stress conditions, and that the two tissues have the similar antioxidant mechanisms.

To summarize, this studies finds that: 1) AANAT2 expression, MT-R1 mRNA and protein expression, and melatonin concentrations in the pineal gland and plasma are controlled by light, 2) gut melatonin differs from pineal

gland melatonin, possibly because it possesses an autonomous circadian rhythm, which is not affected by light, 3) the function of melatonin as an antioxidant does not differ across photosensitive (pineal gland) and non-photosensitive tissue (gut). However, because so few studies have focused on gut melatonin, further research is required before we can fully understand the function of gut melatonin and the mechanisms involved in its secretion.



Acknowledgements

학부 3학년 때부터 해양분자환경생리학 연구실에 들어와 대학원을 진학하여 석사과정을 2년 만에 기쁜 마음으로 마치게 되었습니다. 그동안 아낌없는 조언과 가르침, 힘을 주신 분들께 석사학위를 작성하면서 감사의 마음을 전하고자 합니다.

먼저 과학적 지식을 가르쳐 주시고 부족한 저에게 진심어린 격려와 지적, 관심을 보내주신 지도교수님이신 최철영 교수님께 진심으로 감사를 드립니다. 그리고 대학원 과정 동안 제게 전공분야의 학문적 지식을 알려주신 박인석 교수님과 조성환 교수님, 학부 기간동안 저의 학문의 기초를 쌓게 도움을 주신 안종웅 교수님, 임선영 교수님, 서영완 교수님께 깊은 감사를 드립니다. 논문 심사를 맡아주신 최영웅 박사님께 감사의 말씀을 드립니다.

해양분자환경생리학 연구실에서 적응하게 도와주시고, 실험과 논문 작성을 하면서 큰 힘이 되어주신 나나 누나와 영재형, 그리고 학부 때부터 연구실 생활을 하면서 같이 기쁨과 어려움을 나눈 진아, 아무말 없이 도와 주신 서진이 누나, 든든한 태환, 종렬, 현준, 윤섭, 설혜에게도 감사의 뜻을 전합니다. 그리고 힘든 대학원 생활에 가장 큰 힘, 활력소가 되어준 나의 친구들 <채길, 상호, 윤근, 성업, 성용, 태호>에게도 고마움을 전합니다. 마지막으로 항상 저를 기도로, 좋은 말씀으로 응원하여 주시고, 언제나 저의 주위에서 부족함 찾아서 채워주시는 아빠, 엄마, 지호형 정말 감사합니다. 그리고 사랑합니다.

논문이 완성되기까지 도와주신 모든 분들께 감사의 말씀을 올리며, 언제나 성실하고, 초심을 잃지 않는 지용이가 되겠습니다. 도움을 주셔서 감사합니다.

References

- Benyassi, A., Schwartz, C., Coon, S.L., Klein, D.C., Falcón, J., 2000. Melatonin synthesis: Arylalkylamine N-Acetyltransferases in trout retina and pineal organ are different. *Neuroreport* 11, 255–258.
- Besseau, L., Benyassi, A., Møler, M., Coon, S.L., Weller, J.L., Boeuf, G., Klein, D.C., Falcón, J., 2006. Melatonin pathway: breaking the ‘high-at-night’ rule in trout retina. *Exp. Eye. Res.* 82, 620–627.
- Bubenik, G.A., 1980. Immunohistological localization of melatonin in the salivary glands of the rat. *Adv. Biosci.* 29, 95–112.
- Bubenik, G.A., 2002. Gastrointestinal melatonin: localization, function, and clinical relevance. *Digest. Dis. Sic.* 47, 2336–2348.
- Bubenik, G.A., Brown, G.M., 1997. Pinealectomy reduces melatonin levels in the serum but not in the gastrointestinal tract of rats. *Biol. Signals* 6, 40–4.
- Bubenik, G.A., Pang, S.F., 1997. Melatonin levels in the gastrointestinal tissues of fish, amphibians and a reptile. *Gen. Comp. Endocrinol.* 106, 415–419.
- Bustin, S.A., Benes, V., Garson, J.A., Hellems, J., Huggett, J., Kubista, M., Mueller, R., Nolan, T., Pfaffl, M.W., Shipley, G.L., Vandesompele, J., Wittwer, C.T., 2009. The MIQE guidelines: Minimum Information for Publication of Quantitative Real-Time PCR Experiments. *Clin. Chem.* 55, 611–622.
- Choi, J.Y., Song, J.A., Shin, H.S., Choi, Y.J., Kim, B-S., Yun, S.G., Choi, C.Y., 2014. Effect of LED light spectra on circadian rhythms in goldfish *Carassius auratus*: expression profiles following thermal stress. *Biol.*

- Rhythm Res. 45, 895–908.
- Coon, S.L., Bégay, V., Deurloo, D., Falcón, J., Klein, D.C., 1999. Two arylalkylamine N-acetyltransferase genes mediate melatonin synthesis in fish. *J. Biol. Chem.* 274, 9076–9082.
- Falcón, J., Besseau, L., Sauzet, S., Boeuf, G., 2007. Melatonin and neuroendocrine regulations in fish. *J. Soc. Biol.* 201, 21–29.
- Gothilf, Y., Coon, S.L., Toyama, R., Chitnis, A., Namboodiri, M.A.A., Klein, D.C., 1999. Zebrafish serotonin N-acetyltransferase-2: marker for development of pineal photoreceptors and circadian clock function. *Endocrinology* 140, 4895–4903.
- Iuvone, P. M., Toisni, G., Pozdeyev, N., Haque, R., Klein, D.C., Chaurasia, S.S., 2005. Circadian clocks, clock networks, arylalkylamine N-acetyltransferase, and melatonin in the retina. *Prog. Ret. Eye. Res.* 24, 433–456.
- Kim, N.N., Choi, Y.J., Shin, H.S., Lim, J.R., Han, J.M., Cho, J.H., Lee, J., Kil, G.S., Choi, C.Y., 2014. The effect of LED light spectra on antioxidant system by thermal stress in goldfish, *Carassius auratus*. *Mol. Cell Toxicol.* 10, 47–58.
- Klein, D.C., 2007. Arylalkylamine N-acetyltransferase: “the timenzyme”. *J. Biol. Chem.* 282, 4233–4237.
- Klein, D.C., Coon, S.L., Roseboom, P.H., Weller, J.L., Bernard, M., Gastel, J.A., Zatz, M., Iuvone, P.M., Rodriguez, I.R., Begay, V., Falcon, J., Cahill, G.M., Cassone, V.M., Baler, R., 1997. The melatonin rhythm-generating enzyme: molecular regulation of serotonin N-acetyltransferase in the pineal gland. *Recent Prog. Horm. Res.* 52, 307-358.

- Klein, D.C., Ganguly, S., Coon, S., Weller, J.L., Obsil, T., Hickman, A., Dyda, F., 2002. 14-3-3 Proteins and photoneuroendocrine transduction: role in controlling the daily rhythm in melatonin. *Biochem. Soc. Trans.* 30, 365–373.
- Martinez-Chavez, C.C., Migaud, H. 2009. Retinal light input is required to sustain plasma melatonin rhythms in Nile tilapia *Oreochromis niloticus*. *Brain Res.* 1269, 61–67.
- Migaud, H, Taylor, J.F., Taranger, G.L., Davie, A., Cerda-Reverter, J.M., Carrillo, M., Hansen, T., Bromage, N.B. 2006. A comparative ex vivo and in vivo study of day and night perception in teleosts species using the melatonin rhythm. *J. Pineal Res.* 41, 42–52.
- Mukherjee, S., Moniruzzaman, M., Maitra, S.K., 2014a. Daily and seasonal profiles of gut melatonin and their temporal relationship with pineal and serum melatonin in carp *Catla catla* under natural photo-thermal conditions. *Biol. Rhythm Res.* 45, 301–315.
- Mukherjee, S., Moniruzzaman, M., Maitra, S.K., 2014b. Impact of artificial lighting conditions on the diurnal profiles of gut melatonin in a surface dwelling carp (*Catla catla*) *Biol. Rhythm Res.* 45, 831–848.
- Nakano, T., Kameda, M., Shoji, Y., Hayashi, S., Yamaguchi, T., Sato, M., 2014. Effect of severe environmental thermal stress on redox state in salmon. *Redox. Biol.* 2, 772–776.
- Park, M.S., Shin, H.S., Kim, N.N., Lee, J.H., Kil, G-S., Choi, C.Y., 2013. Effects of LED spectral sensitivity on circadian rhythm-related genes in the yellowtail clownfish, *Amphiprion clarkii*. *Anim. Cells Syst.* 17, 99–105.
- Reiter, R.J., Tan, D.X., Osuna, C., Gitto, E., 2000. Actions of melatonin in

- the reduction of oxidative stress. *J. Biomed. Sci.* 7, 444–58.
- Reppert, S.M., Weaver, D.R., Godson, C., 1996. Melatonin receptors step into the light: cloning and classification of subtypes. *Trends Pharmacol. Sci.* 17, 100–102.
- Underwood, H., Goldman, B.D., 1987. Vertebrate circadian photoperiodic system: role of melatonin and the pineal gland. *J. Biol. Rhythms* 2, 279–315.
- Velarde, E., Cerdá-Reverter, J.D., Alonso-Gómez, A.R., Sánchez, E., Isorna, E., Delgado, M. J., 2010. Melatonin-synthesizing enzymes in pineal, retina, liver, and gut of the goldfish (*Carassius*): mRNA expression pattern and regulation of daily rhythms by lighting conditions. *Chronobiol. Int.* 27, 1178–1201.
- Wu, Y.H., Swaab, D.F., 2005. The human pineal gland and melatonin in aging and Alzheimer's disease. *J. Pineal Res.* 38, 145–152.
- Yamazaki, S., Goto, M., Menaker, M., 1999. No evidence for extraocular photoreceptors in the circadian system of the Syrian hamster. *J. Biol. Rhythms* 14, 197–201.
- Yoshikawa, T., Oishi, T., 1998. Extraretinal photoreception and circadian systems in nonmammalian vertebrates. *Comp. Biochem. Physiol. B* 119, 65–72.